

Brown Spot Severity and Yield of Soybeans Regenerated from Calli Resistant to a Host-Specific Pathotoxin Produced by *Septoria glycines*

G. B. Lee, Former Graduate Student, Department of Crop Sciences, University of Illinois at Urbana-Champaign (UIUC); G. L. Hartman, USDA-ARS, Crop Protection Research Unit and Department of Crop Sciences, UIUC, Urbana, IL 61801; and S. M. Lim, Department of Plant Pathology, University of Arkansas, Fayetteville 72701

ABSTRACT

Lee, G. B., Hartman, G. L., and Lim, S. M. 1996. Brown spot severity and yield of soybeans regenerated from calli resistant to a host-specific pathotoxin produced by *Septoria glycines*. Plant Dis. 80:408-413.

Regenerated soybean lines from the R_3 to R_7 generations were field-evaluated for their reactions to *Septoria glycines*, the brown spot pathogen, from 1991 to 1994. The regenerated lines from cvs. BSR201, Fayette, and L86P-1615 were selected from calli resistant to a host-specific pathotoxin produced by *S. glycines*. Selected resistant plants to the pathogen from the R_3 generation produced R_4 progeny that were resistant, intermediate and susceptible in their reaction to *S. glycines* in 1991. In 1992, 1993, and 1994 among the R_3 to R_7 generations, brown spot reaction was only intermediate and susceptible. F_3 families obtained from a cross between R_3 regenerants and BSR201 had low heritability (23%) for resistance to *S. glycines*. Lines in the R_5 or R_6 generation, obtained originally from plants regenerated from calli of BSR201 that had intermediate and susceptible reactions to *S. glycines* in 1992, and five commercial soybean cultivars were evaluated for brown spot resistance, maturity, and yield in 1993 and 1994. Area under the disease progress curve (AUDPC), maturity, and yield varied among entries. The 10 lines selected with an intermediate reaction had lower AUDPC, matured later, and had higher yields than the nine susceptible lines. Three regenerated lines, 2728, 2733, and 2734, had significantly ($P = 0.05$) lower AUDPC, later maturity, and higher yields than the other regenerated lines. AUDPC was significantly ($P = 0.01$) negatively correlated to yield ($r = -0.29$), to plant height ($r = -0.74$) and number of nodes ($r = -0.39$); but positively correlated to pods having 0 seed ($r = 0.44$) and two seeds ($r = 0.49$).

Brown spot of soybeans, *Glycine max* (L.) Merr., caused by *Septoria glycines* Hemmi, is the most prevalent foliar disease of soybeans in Illinois (4,19). Brown spot causes significant yield losses to soybeans (12,18,26,28). Commercial cultivars, plant introductions, and wild *Glycine* spp. have been tested for resistance to *S. glycines* (11,13,27,29). Resistance to brown spot was detected in two accessions of *G. clandestina* Wendl. and one accession of *G.*

tabacina (Labill.) Benth. (13). Some degree of resistance was detected in 19 lines of *G. max* in Russia (29). Immunity or single gene resistance in soybeans to *S. glycines* has not been reported.

Rapidly spreading leaf chlorosis is a symptom of brown spot that may be

caused by a diffusible substance produced by the fungus (14). A pathotoxin from fungal culture filtrates was purified and partially characterized (21). The toxin was host-specific and caused brown spot symptoms on soybean cotyledons and leaves (21). This suggested that brown spot symptoms may be closely related to the host-specific pathotoxin produced by *S. glycines*.

Resistance to brown spot has been difficult to find by conventional screening methods. One thousand soybean plant introductions were field screened and all were classified as susceptible to *S. glycines*, with >75% leaf area diseased (11). For diseases like brown spot, for which sources of resistance have been difficult to find, somaclonal variation and in vitro selection with pathotoxic culture filtrates have been used for developing disease-resistant plants (1,6-8,22). Soybean plants with resistance to *S. glycines* were selected by means of a pathotoxic culture filtrate of *S. glycines* from cultured cells of cultivars susceptible to brown spot (22). The progenies of these regenerated plants had some level of brown spot resistance in the field (22). In this study, the progenies of those regenerated plants were tested for their brown spot reaction in the R_3 to R_7 generations. In addition, selections from the

Table 1. Generation advancement of soybean lines obtained from calli of cvs. BSR201, Fayette, and L86P-1615, selected for resistance to a pathotoxin produced by *Septoria glycines*

Year ^a	BSR201		Fayette		L86P-1615	
	Generation	Plants or rows	Generation	Plants or rows	Generation	Plants or rows
1988	R_0^b	5 ^c	R_0	1 ^c	R_0	2 ^c
1989	R_1	— ^d	R_1	—	R_1	—
1990	R_2	832 ^c	R_2	—	—	—
1991	R_3	1,651 ^c	R_3	—	R_2	—
1992	R_4	— ^d	—	—	—	—
1992	R_4 and R_5	1,134 ^c	R_4	136 ^c	R_3	936 ^c
1992	R_5	1,046 ^c	—	—	—	—
1992	F_2 and F_3	317 ^e	—	—	—	—
1993	R_5 and R_6	460 ^c	R_5	16 ^e	R_4	34 ^e
1993	F_3	100 ^e	—	—	—	—
1993	F_4	1,549 ^c	—	—	—	—
1994	R_6 and R_7	96 ^c	R_6	14 ^e	R_5	15 ^e
1994	F_5	31 ^c	—	—	—	—
1994	F_5	20 ^e	—	—	—	—

^a Regenerated lines in 1988 to 1990 were previously advanced (22).

^b Generation number.

^c Number of plants evaluated following field inoculation.

^d Unknown number of plants advanced.

^e Number of rows evaluated following field inoculation.

Corresponding author: G. L. Hartman
E-mail: ghartman@uiuc.edu

Present address of first author: Crop Experiment Station (RDA), 209 Sotundong, Suwon, Korea, 441-100.

Supported in part by the Illinois Soybean Operating Board.

Mention of a trademark or proprietary product does not constitute a guarantee or warranty of the product by the U.S. Department of Agriculture and does not imply its approval to the exclusion of other products that may also be suitable.

Accepted for publication 28 December 1995.

Publication no. D-1996-0207-04R

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 1996.

R₅ and R₆ generations were evaluated for yield and brown spot resistance.

MATERIALS AND METHODS

Source of regenerants. The R₃ to R₇ lines used were advanced from R₃ lines reported to have resistance to *S. glycines*, based on reaction of calli to the toxin produced by this fungus (22). The R₃ to R₇ progeny of regenerated lines and F₂ to F₅ progeny of crosses between regenerated lines from cv. BSR201 and BSR201 (Table 1) were tested for resistance to *S. glycines* in the field.

Evaluation of resistance. From 1991 to 1994, seeds were collected from plants that had a resistant or intermediate reaction to brown spot. Seeds from each plant were sown in a single row the following year. The cvs. BSR201, Fayette, and L86P-1615 were used as checks each year. Seeds were planted in 2.4-m rows on 28 May 1991, 19 May 1992, 17 May 1993, and 20 May 1994. Plants at growth stages V2 to V6 (5) were inoculated with *S. glycines* by means of a pressurized sprayer (5.6 kg/cm²) on 28 June, and 2 and 15 July 1991; 15 and 29 June and 10 July 1992; 22 June 1993; and 27 June and 19 July 1994. Inoculum was produced by growing *S. glycines* on potato dextrose agar (PDA) for 2 weeks at 24 ± 2°C. Cultures were macerated in a blender with tap water for 1 to 2 min and adjusted to 10⁶ conidia/ml. Disease severity was rated using a modified Horsfall-Barratt scale based on visual observations of the proportion of leaf area diseased (9, 17). Disease severity and plant growth stage were evaluated weekly for 5 to 7 weeks. Reactions of individual regenerants or plant rows were classified into three categories based on percentage of leaf area diseased: resistance = 0 to 6%; intermediate = 7 to 25%; and susceptible = >25%.

Heritability test. R₃ plants from BSR201 calli that had resistance to *S. glycines* were pollinated with BSR201 in the greenhouse in 1990. Seeds of the F₁ generation (R₃F₁) were advanced to the F₂ generation (R₃F₂) in the field in 1992. Eighty-three F₃ families (R₃F₃) were randomly selected and seeds were planted along with BSR201 at the rate of 25 seeds per 75-cm row on 17 May 1993. The experiment was arranged in randomized complete block design with two replications. Inoculation method, time of inoculations, and rating scale were the same as previously described for the resistance evaluation in 1993. A total of 1,549 plants were individually evaluated for brown spot severity at growth stage R6. Severity data for individual plants in the 83 families were analyzed by analysis of variance. Genetic variability, phenotypic variability, and heritability were estimated as follows: $s^2G = [(s^2E + r s^2G) - s^2E]/r$, $s^2P = s^2G + s^2E$, and $h^2 = s^2G / s^2P$, respectively, where s^2G is genetic variability, s^2E is environmental variability, s^2P is phenotypic variability, and h^2 is heritability (16).

Evaluation of resistance and yield. Based on field evaluations in 1992, seeds from 10 plants that had an intermediate reaction to *S. glycines* and nine plants that were susceptible from the R₅ and R₆ generations from BSR201 calli were selected and planted along with five commercial cvs. (Asgrow A2943, BSR201, Burlison, Kenwood, and Sturdy) in 3.6-m-long two-row plots, with a distance between rows of 76 cm, in a continuous soybean cropping field on 19 May 1993. After germination, rows were trimmed to a length of 2.4 m. All plots were inoculated with a suspension of 10⁶ conidia/ml concentration by means of a pressurized sprayer (5.6 kg/cm²). Preparation of inoculum was the

same as described previously. Entries were replicated three times in a randomized complete block design. The percentage of disease severity and the growth stages were visually estimated on a whole plot basis weekly from 7 July to 1 September. The area under disease progress curve (AUDPC) was calculated as: $\sum N_i = 1 ((R_i + R_{i+1})/2)(t_i + 1 - t_{i+1})$, where t_i = day of evaluation i , R_i = percentage of disease severity at evaluation i , and i = one to eight evaluations (20). After maturity all plants in rows were harvested and dried at 37°C for 72 h to 8% moisture. Yields were adjusted to 13% moisture.

In 1994, the same 24 entries as in 1993 were planted in 4.8-m long four-row plots in a soybean-corn rotation field and in a continuous soybean cropping field. After seed germination, plant rows were trimmed to 3.6 m. A split-plot randomized complete block design was used with three replications. The inoculated and noninoculated plots were arranged as main plots and the 24 lines/cultivars were arranged as subplots. Plants were inoculated with a suspension of 10⁶ conidia/ml concentration as described previously. Noninoculated plots were sprayed with five applications of benomyl (Benlate 50% WP formulation: methyl 1-[butylcarbamoyl]-2-benzimidazolecarbamate) applied at 1.1 kg/ha using 280 liters of water per ha. Disease severity was rated weekly on whole plots from 21 July to 31 August using the modified Horsfall-Barratt scale. AUDPC was calculated as described previously. At maturity, height and number of nodes on the main stem were recorded for 10 plants from each plot. Twenty plants in each plot for five regenerated lines (three best lines and two worst lines for AUDPC based on 1993 data) and BSR201 were sampled after maturity. All pods were removed and separated to determine the number of seeds per pod. The center two rows of all plots were harvested, dried, and weighed for yield as described previously. Data

Table 2. Field evaluation of brown spot in different generations of soybean lines regenerated from calli that were selected for resistance to a pathotoxin produced by *Septoria glycines*

Year	Generation and source ^a	Plants evaluated	Rows evaluated	Reaction (percentage) ^b		
				Resistant	Intermediate	Susceptible
1991	R ₄ P1	1,134	45	148 ^c (13)	890 ^c (78)	96 ^c (9)
	BSR201	28	1	0	25 ^c (89)	3 ^c (11)
1992	R ₃ P2	936	36	0	82 ^c (9)	854 ^c (91)
	R ₄ P1	291	15	0	14 ^c (5)	277 ^c (95)
	R ₄ P3	136	5	0	14 ^c (5)	122 ^c (90)
	R ₅ P1	755	80	0	9 ^c (1)	746 ^c (99)
	F ₃ P1	330	18	0	11 ^c (3)	319 ^c (97)
	BSR201	65	2	0	0	65 ^c (100)
	R ₄ P2	0	34	0	2 ^d (6)	32 ^d (94)
1993	R ₅ P1	0	32	0	2 ^d (6)	30 ^d (94)
	R ₅ P3	0	16	0	5 ^d (31)	11 ^d (69)
	R ₆ P1	0	68	0	3 ^d (4)	65 ^d (96)
	Check cultivars ^e	0	15	0	0	15 ^d (100)

^a R₃ to R₆ indicates the number of selfed generations from organogenic cultures. F₃ indicates generation of a cross between a regenerated line with BSR201. P1 indicates calli were derived from BSR201. P2 indicates calli were derived from L86P-1615. P3 indicates calli were derived from Fayette.

^b Resistant = 1 to 6%; intermediate = 7 to 25%; susceptible = > 25% of leaf area affected.

^c Total number of plants evaluated.

^d Total number of rows evaluated.

^e Cvs. BSR201, Fayette, and L86P-1615.




Fig. 1. Brown spot progress curves for soybean cv. BSR201 and soybean lines in the R₅ (from calli of L86P-1615), R₆ (from calli of Fayette), and R₇ (from calli of BSR201) generations after inoculation with *Septoria glycines* in the field in 1994.

Table 3. Field evaluation of brown spot on soybean cv. BSR201 regenerated lines that were selected one generation earlier based on their reaction to *Septoria glycines*^a

Reaction ^b	Generation ^c	No. of rows	Reaction (percentage)	
			Intermediate	Susceptible
Intermediate	R ₅	293	42 ^d (14)	251 (86)
Intermediate	F ₄	51	3 (4)	50 (96)
Intermediate	F ₅	4	4 (100)	0 (0)
Susceptible	R ₅	24	0 (0)	24 (100)
Susceptible	F ₄	45	0 (0)	45 (100)
Susceptible	F ₅	6	0 (0)	6 (100)
Susceptible	BSR201	14	0 (0)	14 (100)

^a Disease severity based on ratings at growth stages R₄ to R₅ from 1992 to 1994.

^b Intermediate = 7 to 25% and susceptible = >25% leaf area diseased.

^c R₅ is the fifth-selfed generation. F₄ and F₅ are the fourth- and fifth-selfed generations from R₃ and R₂ regenerants, respectively, crossed to BSR201. R₅, F₄, and F₅ generations were evaluated in 1992, 1993, and 1994, respectively.

^d Total number of rows evaluated.

Table 4. Area under disease progress curve (AUDPC) and relative rank, maturity date, and yield and relative rank for 19 soybean lines regenerated from organogenic cultures of cv. BSR201 and five commercial cultivars inoculated with *Septoria glycines* in the field in 1993

Lines/cultivars	AUDPC		Maturity date	Yield	
	Value	Rank		Kg/ha	Rank
Intermediate ^a					
2322	1599	5	Sept. 14	3,172	7
2335	1630	7	Sept. 13	2,972	17
2354	1605	6	Sept. 14	3,114	12
2387	1681	12	Sept. 13	3,015	14
2681	1649	8	Sept. 14	2,991	15
2700	1652	9	Sept. 14	3,156	9
2709	1574	4	Sept. 14	3,123	11
2728	1292	3	Sept. 19	3,185	6
2733	1290	2	Sept. 18	3,511	2
2734	1242	1	Sept. 20	3,113	13
Susceptible ^b					
2281	1735	17	Sept. 12	3,159	8
2284	1763	19	Sept. 11	2,811	23
2325	1811	24	Sept. 11	2,836	22
2330	1786	22	Sept. 10	2,954	18
2442	1713	15	Sept. 10	2,793	24
2447	1725	16	Sept. 11	2,977	16
2455	1796	23	Sept. 10	2,858	21
2692	1696	14	Sept. 12	2,932	19
2699	1658	10	Sept. 13	3,134	10
Check cultivars					
ASG2943	1666	11	Sept. 18	3,412	4
BSR201	1739	18	Sept. 14	2,928	20
Burlison	1695	13	Sept. 17	3,450	3
Kenwood	1778	21	Sept. 14	3,672	1
Sturdy	1774	20	Sept. 10	3,203	5
FLSD ^c	90			220	
Means					
Intermediate/10 lines	1521		Sept. 15	3,135	
Susceptible/9 lines	1742		Sept. 11	2,939	
Check cultivars	1730		Sept. 15	3,333	
Three lowest AUDPC lines ^d	1274		Sept. 19	3,268	
Comparisons					
Intermediate vs. susceptible	0.01 ^e		0.01	0.01	
Intermediate vs. checks	0.01		NS ^f	0.01	
Susceptible vs. checks	NS ^f		0.01	0.01	
Three lowest AUDPC lines vs. BSR201	0.01		0.01	0.01	

^a Intermediate = 10 lines with intermediate reaction from progenies of R₅ and R₆ generations.

^b Susceptible = nine lines with susceptible reaction from progenies of R₅ and R₆ generations.

^c Fisher's protected least significant difference ($P = 0.05$).

^d Three lowest AUDPC lines = 2728, 2733, and 2734.

^e Means differ significantly at $P = 0.01$.

^f Not significant.

were analyzed by analysis of variance. Entries were divided into two groups, inoculated and noninoculated plots, and the means were separated by Fisher's protected least significant difference and group mean comparison.

RESULTS

Evaluation of resistance. In 1991, 148 of 1,134 R₄ generation plants were classified as resistant, 890 as intermediate, and 96 along with BSR201 as susceptible (Table 2). In 1992, 1 to 10% of the regenerants had an intermediate reaction, and all the rest were susceptible, as was BSR201 (Table 2). In 1993, no plants were rated resistant (Table 2). Only 4 to 31% of the 150 rows in the different generations had an intermediate reaction, and all the rest including the check cultivars were susceptible (Table 2). The disease classification of regenerants varied based on the source of parental calli. For example, regenerants in the R₄ and R₅ generations from calli of Fayette and L86P-1615 had more plants with an intermediate reaction (10 and 9%, respectively) than those from BSR201 in the R₄ and R₆ generations in 1992 and 1993 (Table 2).

BSR201 and three lines selected based on an intermediate reaction that were derived from calli of cvs. BSR201, Fayette, and L86P-1615 had slow disease development in 1994 until 38 days after inoculation, at which time brown spot greatly increased (after growth stage R₅) (Fig. 1). Disease progress curves for BSR201 and the three lines were similar until growth stage R₅. After growth stage R₅ all selected lines had significantly ($P = 0.01$) lower disease severity than BSR201 (Fig. 1).

The R₅, F₄, and F₅ lines selected for susceptibility in the previous year were susceptible the next year. R₅ and F₄ lines selected for their intermediate reaction in the previous year varied from intermediate to susceptible, while F₅ lines from the intermediate reaction in the previous year were intermediate the next year (Table 3).

Heritability test. The mean square variance for 83 R₃F₃ families was 27 and the error mean square variance was 17. The genetic variance was 5, environmental variance was 17, and phenotypic variance was 22. Heritability of disease resistance in R₂F₃ families was 23% of the total genetic variance.

Evaluation of resistance and yield. In 1993, AUDPC varied among the entries from 1,242 to 1,811 (Table 4). Mean AUDPC of the 10 intermediate lines was significantly ($P = 0.01$) less than the mean AUDPC of the nine susceptible lines and the five check cultivars. Maturity also varied among the entries from 10 to 20 September (Table 4). The intermediate lines matured later than the susceptible lines, but had a maturity similar to that of the five check cultivars. Yield also varied

among the entries from 2,793 to 3,672 kg/ha (Table 4). Average yields of the 10 intermediate lines (3,135 kg/ha) was significantly ($P = 0.01$) higher than the average yield of 2,939 kg/ha for the nine susceptible lines (Table 4). Yields of three lowest AUDPC lines were significantly ($P = 0.01$) higher than that of BSR201.

When benomyl was used to reduce brown spot severity, maturity and yield varied among entries in 1994 (Table 5). The AUDPC of the 10 intermediate lines was similar to that of the nine susceptible lines and the five check cultivars. Days to maturity of the 10 intermediate lines were delayed by 6 days compared with the susceptible lines and BSR201. The 10 intermediate lines had a yield similar to that of the nine susceptible lines and the five check cultivars. BSR201 ranked 14th for yield and had a yield similar to that of the three lowest AUDPC lines.

In the inoculated trial in 1994, mean AUDPC of the 10 intermediate lines was significantly ($P = 0.01$) lower than mean AUDPC of the nine susceptible lines and mean AUDPC of the five check cultivars (Table 6). BSR201 ranked 18th for AUDPC among 24 entries. The 10 intermediate lines matured later than the nine susceptible lines and BSR201. Three lines (2728, 2733, and 2734) had significantly ($P = 0.01$) lowest AUDPC compared with the other intermediate and susceptible lines, and check cultivars. Yields of the 10 intermediate lines were significantly ($P = 0.01$) higher than those of the nine susceptible lines and BSR201, but were similar to the five check cultivars.

When comparing AUDPC between the inoculated and fungicide-protected plots, the AUDPC in the inoculated plots for the 10 intermediate lines was less than that of the nine susceptible lines and the five check cultivars (Table 7). Days to maturity increased for plants in fungicide-protected plots compared with inoculated plots with BSR201 having the greatest difference. The yield loss of the 10 intermediate lines was lower than that of the nine susceptible lines and the five check cultivars. BSR201 had 16% less yield in the inoculated trial than in the protected trial.

AUDPC was significantly ($P = 0.01$) negatively correlated to yield ($r = -0.29$), to plant height ($r = -0.74$) and number of nodes ($r = -0.39$); but positively correlated to pods having 0 seed ($r = 0.44$) and two seeds ($r = 0.49$).

DISCUSSION

Pathotoxins have been used for the selection of in vitro resistance for alfalfa, corn, potato, soybean, and tobacco (1,6-8, 22,24). In vitro selection and regeneration of resistant soybean plants to *S. glycines* in the R_2 and R_3 generations were reported under field screening (22). Regenerated plants when inoculated with *S. glycines* at growth stage V3 had no brown spot until

growth stage R6. All resistant plants were shorter, and varied in maturity and sterility (22). Selection of both resistant and susceptible regenerated plants produced progeny that varied in resistance to *S. glycines* (22). In our study, the regenerants from the R_3 to R_7 generations differed from the R_2 and R_3 generations in resistance to brown spot. Brown spot resistance was not observed in the R_3 to R_7 generations from 1992 to 1994 although a few plants were classified as resistant in 1991. The growing season in 1991 was drier than that of other years, which reduced the severity of brown spot. Lines susceptible to brown spot produced susceptible progeny the next year whereas lines intermediate to brown spot produced progenies that varied from intermediate to susceptible. Variations for other characteristics, including plant height, maturity, and sterility, were

detected only from calli of immature embryo selections in a limited number of families. The level of brown spot resistance reported in the R_2 and R_3 generations derived from immature organogenesis (22) was not maintained in advanced generations.

In order for in vitro selection to be successful, traits should be heritable. This has not, however, always been true, since resistant variants expressed in culture were not always resistant in regenerated plants (3, 25). In addition, other traits have been shown to be lost upon regeneration (2,15). In our study, most of the selected lines in the R_7 generation were susceptible to *S. glycines*. Also a few of the selected lines from calli of immature seeds of BSR201 were not stable for resistance to *S. glycines*, or for maturity and sterility. Lines regenerated from different calli may differ

Table 5. Area under disease progress curve (AUDPC) and relative rank, days to maturity, and yield and relative rank of 19 soybean lines regenerated from organogenic cultures of cv. BSR201 and five commercial cultivars following protection with benomyl in the field in 1994

Lines or comparison	AUDPC		Maturity date	Yield	
	Value	Rank		Kg/ha	Rank
Intermediate ^a					
2322	27	5	Sept. 12	2,544	23
2335	35	18	Sept. 12	2,993	10
2354	37	19	Sept. 14	3,165	4
2387	38	21	Sept. 13	2,906	13
2681	33	10	Sept. 16	3,108	7
2700	26	2	Sept. 13	2,815	17
2709	39	22	Sept. 15	3,119	6
2728	24	1	Sept. 16	2,695	21
2733	34	13	Sept. 19	3,163	5
2734	33	11	Sept. 19	3,460	1
Susceptible ^b					
2281	26	3	Sept. 11	2,510	24
2284	35	16	Sept. 11	2,947	12
2325	35	17	Sept. 12	2,818	16
2330	34	12	Sept. 14	2,876	15
2442	28	6	Sept. 8	2,559	22
2447	38	20	Sept. 10	2,789	18
2455	43	24	Sept. 9	2,742	20
2692	35	15	Sept. 13	3,073	8
2699	34	14	Sept. 13	3,003	9
Check cultivars					
ASG2943	31	8	Sept. 17	3,377	2
BSR201	40	23	Sept. 13	2,880	14
Burlison	29	7	Sept. 14	2,773	19
Kenwood	27	4	Sept. 12	3,359	3
Sturdy	31	9	Sept. 9	2,984	11
FLSD ^c	NS ^d			372	
Means					
Intermediate/10 lines	32		Sept. 15	2997	
Susceptible/9 lines	34		Sept. 11	2813	
Check cultivars	32		Sept. 13	3075	
Three lowest AUDPC lines ^e	30		Sept. 18	3106	
Comparisons					
Intermediate vs. susceptible	NS		0.01	NS	
Intermediate vs. check cultivars	NS		0.01	NS	
Susceptible vs. check cultivars	NS		0.05	0.05	
Three lowest AUDPC lines vs. BSR201	0.05 ^f		0.01	NS	

^a Intermediate = 10 lines with intermediate reaction from progenies of R_5 and R_6 generations.

^b Susceptible = nine lines with susceptible reaction from progenies of R_5 and R_6 generations.

^c Fisher's protected least significant difference ($P = 0.05$).

^d Not significant.

^e Three lowest AUDPC lines = 2728, 2733, and 2734.

^f Means differ significantly at $P = 0.05$.

Table 6. Area under disease progress curve (AUDPC) and relative rank, days to maturity, yield and relative rank for 19 soybean lines regenerated from organogenic cultures of cv. BSR201 and five commercial cultivars following inoculation with *Septoria glycines* in the field in 1994

Lines or comparison	AUDPC		Maturity date	Yield	
	Value	Rank		Kg/ha	Rank
Intermediate^a					
2322	833	14	Sept. 7	2,388	13
2335	860	15	Sept. 6	2,253	19
2354	791	9	Sept. 8	2,278	18
2387	788	8	Sept. 7	2,481	10
2681	751	6	Sept. 8	2,506	8
2700	749	5	Sept. 9	2,553	7
2709	732	4	Sept. 9	2,498	9
2728	570	1	Sept. 12	2,868	2
2733	585	3	Sept. 13	2,725	4
2734	576	2	Sept. 14	2,890	1
Susceptible^b					
2281	895	17	Sept. 4	2,425	11
2284	902	20	Sept. 4	2,106	23
2325	891	16	Sept. 4	2,239	20
2330	910	21	Sept. 3	2,346	15
2442	953	22	Sept. 10	2,116	22
2447	897	19	Sept. 8	2,282	17
2455	973	24	Sept. 3	2,103	24
2692	779	7	Sept. 7	2,354	14
2699	812	12	Sept. 7	2,336	16
Check cultivars					
ASG2943	821	13	Sept. 11	2,641	6
BSR201	895	18	Sept. 5	2,420	12
Burlison	810	11	Sept. 11	2,646	5
Kenwood	801	10	Sept. 7	2,795	3
Sturdy	973	23	Aug. 31	2,230	21
FLSD^c	99			306	
Means					
Intermediate/10 lines	723		Sept. 9	2,544	
Susceptible/9 lines	890		Sept. 6	2,256	
Check/5 cultivars	860		Sept. 7	2,546	
Three lowest AUDPC lines ^d	577		Sept. 13	2,827	
Comparisons					
Intermediate vs. susceptible	0.01 ^e		0.01	0.01	
Intermediate vs. check cultivars	0.01		0.01	NS	
Susceptible vs. check cultivars	NS ^f		NS	0.01	
Three lowest AUDPC lines vs. BSR201	0.01		0.01	0.01	

^a Intermediate = 10 lines with intermediate reaction from progenies of R5 and R6 generations.

^b Susceptible = nine lines with susceptible reaction from progenies of R5 and R6 generations.

^c Fisher's protected least significant difference ($P = 0.05$).

^d Three lowest AUDPC lines = 2728, 2733, and 2734.

^e Means differ significantly at $P = 0.01$ and 0.05 . NS = not differ significantly at $P = 0.05$.

^f Not significant.

Table 7. Effect of control of brown spot on disease development, and days to maturity, and yield, of regenerated lines selected in R5 and R6 generations from organogenic cultures resistant to pathotoxin produced by *Septoria glycines*

Character	Lines of comparison	Brown spot severity ^a		Difference
		Low	High	
AUDPC ^b	Average	33	814	+ 781
	10 lines from intermediate	32	723	+ 691
	9 lines from susceptible	34	890	+ 856
	5 check cultivars	32	860	+ 828
	BSR201	40	895	+ 855
	3 lowest AUDPC lines	30	577	+ 547
Maturity date	Average	Sept. 13	Sept. 7	- 6 days
	10 lines from intermediate	Sept. 15	Sept. 9	- 6 days
	9 lines from susceptible	Sept. 11	Sept. 6	- 5 days
	5 check cultivars	Sept. 13	Sept. 7	- 6 days
	BSR201	Sept. 13	Sept. 5	- 8 days
	3 lowest AUDPC lines	Sept. 18	Sept. 13	- 5 days
Yield (kg/ha)	Average	2,944	2,436	- 508 (17) ^a
	10 lines from intermediate	2,997	2,544	- 453 (15)
	9 lines from susceptible	2,813	2,256	- 557 (20)
	5 check cultivars	3,075	2,546	- 529 (17)
	BSR201	2,880	2,420	- 460 (16)
	3 lowest AUDPC lines	3,106	2,827	- 279 (9)

^a Low = benlate treated; high = inoculated with *Septoria glycines*.

^b Area under disease progress curve for brown spot.

genetically and have different backgrounds for disease resistance.

There have been reports of successful in vitro selection for resistance. The selection of T-toxin resistance in tissue culture of corn inbreds resulted in the isolation of many resistant cell lines that were stable over time (6). The regenerated lines from protoplast-derived callus of tobacco were shown to be useful for resistance to wildfire disease (24). In our study, some degree of resistance from three regenerated lines, 2728, 2733, and 2734, obtained from calli of BSR201 were stable in the R₆ and R₇ generations and these also had higher yields. Agronomic traits of these regenerated lines were similar to BSR201. These advanced lines had resistance to brown stem rot (10) as does BSR201 (23). Since some of these lines selected from calli of BSR201 have better resistance to brown spot and have higher yields than BSR201, these advanced lines will be useful as improved germ plasm.

ACKNOWLEDGMENTS

We would like to thank R. L. Warsaw for assistance in the field work.

LITERATURE CITED

- Behnke, M. 1980. General resistance to late blight of *Solanum tuberosum* plants regenerated from callus resistant to culture filtrates of *Phytophthora infestans*. Theor. Appl. Genet. 56:151-152.
- Bins, A., and Meins, F. Jr. 1973. Evidence that habituation of tobacco pith cells for cell division-promoting factors is heritable and potentially reversible. Proc. Natl. Acad. Sci. USA 70:2660-2662.
- Chaleff, R. S., and Keil, R. L. 1981. Genetic and physiological variability among cultured cells and regenerated plants of *Nicotiana tabacum*. Mol. Gen. Genet. 181:254-258.
- Eathington, S. R., Lim, S. M., Nickell, C. D., Pataky, J. K., and Esgar, R. W. 1993. Disease pressure on soybean in Illinois. Plant Dis. 77:1136-1139.
- Fehr, W. R., Caviness, C. E., Burmood, D. T., and Pennington, J. S. 1971. Stage development descriptions of soybean. *Glycine max* (L.) Merr. Crop. Sci. 11:929-931.
- Gengenbach, B. G., Green, C. E., and Donnovan, C. M. 1977. Inheritance of selected pathotoxin resistance in maize plants regenerated from cell cultures. Proc. Natl. Acad. Sci. USA 74:5113-5117.
- Gengenbach, B. G., and Rines, H. W. 1986. Use of phytotoxins in selection of disease resistant mutants in tissue culture. Iowa State J. Res. 60:449-476.
- Hartman, C. L., McCoy, T. J., and Knous, T. R. 1984. Selection of alfalfa (*Medicago sativa*) cell lines and regeneration of plant resistant to the toxin(s) produced by *Fusarium oxysporum* f. sp. *medicaginis*. Plant Sci. Lett. 34:183-184.
- Horsfall, J. G., and Barratt, R. W. 1945. An improved grading system for measuring plant diseases. (Abstr.) Phytopathology 35:655.
- Lee, G. B. 1995. Host range of *Septoria glycines*, quantification of partial resistance, and brown spot evaluation of soybeans regenerated from calli. Ph.D. thesis. University of Illinois at Urbana-Champaign.
- Lim, S. M. 1979. Evaluation of soybean for resistance to *Septoria* brown spot. Plant Dis. Rep. 63:242-245.
- Lim, S. M. 1980. Brown spot severity and

- yield reduction in soybean. *Phytopathology* 70:974-977.
13. Lim, S. M., and Hymowitz, T. 1987. Reactions of perennial wild species of genus *Glycine* to *Septoria glycines*. *Plant Dis.* 71:891-893.
 14. MacNeill, B. H., and Zalasky, H. 1957. Histological study of host-parasite relationships between *Septoria glycines* Hemmi and soybean leaves and pods. *Can. J. Bot.* 35:501-505.
 15. Mok, M. C., Gabelman, W. H., and Skoog, F. 1976. Carotenoid synthesis in tissue cultures of *Daucus carota* L. *J. Am. Soc. Hortic. Sci.* 101:442-449.
 16. Nyquist, W. E. 1991. Estimation of heritability and prediction of selection response in plant populations. *Crit. Rev. Plant Sci.* 10: 235-322.
 17. Park, E. W., and Lim, S. M. 1985. Empirical estimation of asymptotes of disease progress curves and the use of the Richards generalized rate parameters for describing disease progress. *Phytopathology* 75:786-791.
 18. Pataky, J. K., and Lim, S. M. 1981. Effects of row width and plant growth habit on *Septoria* brown spot development and soybean yield. *Phytopathology* 71:1051-1056.
 19. Pataky, J. K., Lim, S. M., Jordan, E. G., and Warsaw, R. L. 1979. Monitoring soybeans for foliar disease. III. Res. 21:3-4.
 20. Shaner, G., and Finney, R. E. 1977. The effect of nitrogen fertilization on the expression of slow mildew resistance in Knox wheat. *Phytopathology* 67:1051-1056.
 21. Song, H. S., Lim, S. M., and Clark, J. M., Jr. 1993. Purification and partial characterization of host-specific pathotoxin from culture filtrates of *Septoria glycines*. *Phytopathology* 83:659-661.
 22. Song, H. S., Lim, S. M., and Widholm, J. M. 1994. Selection and regeneration of soybeans resistant to the pathotoxic culture filtrates of *Septoria glycines*. *Phytopathology* 84:948-951.
 23. Tachibana, H., Bahrenfus, J. B., and Fehr, W. R. 1983. Registration of BSR201 soybean. *Crop Sci.* 23:186.
 24. Thanutong, P., Furusawa, I., and Yamamoto, M. 1983. Resistant tobacco plants from protoplast-derived calluses selected for their resistance to *Pseudomonas* and *Alternaria* toxins. *Theor. Appl. Genet.* 66:209-215.
 25. Widholm, J. 1980. Differential expression of amino acid biosynthetic control isozymes in plants and cultured cells. Pages 157-159 in: *Plant Cell Cultures: Results and Perspectives*. F. Sala and B. Paris, eds. Elsevier, Amsterdam.
 26. Williams, D. J., and Nyvall, R. F. 1980. Leaf infection and yield losses caused by brown spot and bacterial blight diseases of soybean. *Phytopathology* 70:900-902.
 27. Young, L. D., and Ross, J. P. 1978. Resistance evaluation and inheritance of a nonchlorotic response to brown spot of soybean. *Crop. Sci.* 18:1075-1077.
 28. Young, L. D., and Ross, J. P. 1979. Brown spot development and yield response of soybean inoculated with *Septoria glycines* at various growth stages. *Phytopathology* 69:8-11.
 29. Zaostrovnykh, V. I., and Vashchenko, A. P. 1991. Form of soybean resistant to *Septoria*, *Peronospora*, and *Cercospora*. *Sel. Semenovod. (Moscow)* 5:29-30.